

QUALI-PRO®

3-D

Herbicide

FOR USE BY TURF MAINTENANCE PERSONNEL, LANDSCAPING, OR COMMERCIAL APPLICATORS ONLY.

SELECTIVE BROADLEAF WEED CONTROL IN TURFGRASSES INCLUDING USE ON GOLF COURSES AND SOD FARMS. ALSO FOR HIGHWAYS, RIGHTS-OF-WAY, AND OTHER SIMILAR NON-CROP AREAS. TO CONTROL CLOVER, DANDELION, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-P, AND DICAMBA; THE ACTIVE INGREDIENTS USED IN TRIPLET® SELECTIVE HERBICIDE. QUALI-PRO 3-D IS NOT MANUFACTURED OR DISTRIBUTED BY NUFARM AMERICAS, INC.

ACTIVE INGREDIENTS: % BY WT.

*Dimethylamine Salt of 2,4-Dichloro-phenoxyacetic acid	30.56%
**Dimethylamine Salt of (+)-R-2-(2-methyl-4-chlorophenoxy) propionic acid ≠	8.17%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	2.77%
OTHER INGREDIENTS:	<u>58.50%</u>
TOTAL	100.00%

Contains:

*2.44 lbs. 2,4-Dichlorophenoxyacetic acid per gallon or 25.38%.

**0.65 lbs. (+)-R-2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 6.75%.

***0.22 lbs. 3,6-dichloro-o-anisic acid per gallon or 2.30%.

≠CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get into eyes, on skin, or on clothing.

EPA Reg. No. 66222-225

EPA Est. No. 37507-MT-001

For additional first aid, precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

EPA 102510/Rev A

FIRST AID	
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Prosar at 1-877-250-9291 for emergency medical treatment information. NOTE TO PHYSICIAN: If in eyes, specialized ophthalmologic attention may be necessary. If swallowed, probable mucosal damage may contraindicate the use of gastric lavage. There is no specific antidote; treat symptomatically.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate, nitrile rubber, neoprene rubber, or Viton. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Goggles or face shield
- Chemical-resistant gloves, except for applicators using ground boom equipment
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

For containers over 1 gallon and less than 5 gallons in capacity: Persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron.

For containers of 5 gallons or more in capacity: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a nonrefillable pesticide container are emptied, the probe must be rinsed before removal.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets or exceeds the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates and may adversely affect nontarget plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA web site <http://www.epa.gov/essp>.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to

contain spills will help prevent groundwater contamination.

Cleaning of Equipment: When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Do not apply when weather conditions favor drift away from target area.

Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not enter or allow people or pets to enter the treated area until sprays have dried.

PRECAUTIONS AND RESTRICTIONS

Do not enter or allow people or pets to enter the treated area until sprays have dried.

Do not use this product on or near desirable plants including within the drip line of the roots of desirable trees and shrubs, since injury may result. Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product runoff, do not overwater the treated area(s) or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

Avoid broadcast and small spot treatment applications when temperatures exceed 90°F, as turf injury may occur.

Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results, turf should not be mowed for 1 to 2 days before and after application, do not apply if rain is expected within 4 hours after the application, and delay irrigation cycle for 24 hours. Reseed no sooner than 3 to 4 weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not pour spray solutions near these plants.

Chemigation Statement: Do not apply this product through any type of irrigation equipment.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Boom Height

For ground boom applications, do not apply with a nozzle height greater than 4 feet above the crop canopy.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE Standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles. When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas including but not limited to residential areas, bodies of water,

known habitat for nontarget species, nontarget crops within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption. Susceptible crops include but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

ADJUVANT USE: When an adjuvant is to be used with this product, Makhteshim Agan of North America, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

DRIFT CONTROL AND DEFOAMING: For drift control and defoaming, use a product similar to Compadre® at .125% v/v.

FERTILIZER USE: This product can be mixed with some liquid fertilizers or liquid iron metals. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All state and/or federal regulations relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED BY QUALI-PRO 3-D

Bedstraw	Henbit	Purslane
Black Medick	Knottedweed	Ragweed
Buckhorn	Lambsquarters	Sheep Sorrel
Burdock	Lespedeza	Shepherdspurse
Chicory	Mallow	Speedwell
Chickweed	Morningglory	Surge
Clover	Peppergrass	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Dock	Plantain	Wild Lettuce
Ground Ivy	Poison Ivy	Wild Onion
Heal-All	Poison Oak	Yarrow

USE SITES SOD FARMS

Quali-Pro 3-D is intended for use on sod farms. Quali-Pro 3-D provides selective, broadleaf control in warm-season and cool-season turfgrass established for commercial sod production. For application information on sod farms, refer to the tables found in the section of the labeling entitled **ORNAMENTAL AND RECREATIONAL LAWNS AND TURF** for allowable use sites, rates, and comments, but follow the section of the labeling entitled **Use Precautions and Limitations for Sod Farms** in the paragraph below.

Application Schedules for Sod Farms

Apply Quali-Pro 3-D to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to "Floratum" St. Augustinegrass.

Newly Seeded Areas

The application of Quali-Pro 3-D to grass seedlings is suggested after the second mowing.

Newly Sodded, Sprigged, or Plugged Areas

The application of Quali-Pro 3-D to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant bermudagrass, dormant zoysiagrass, and dormant bahiagrass are suggested.

Use Precautions and Limitations for Sod Farms

1. Delay mowing 1 to 2 days before and after the application of this product.
2. Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 24 hours after application.
3. Treated areas may be reseeded 3 to 4 weeks after application.
4. Application is limited to 2 applications per year.
5. Do not exceed a maximum of 4.0 pints Quali-Pro 3-D (1.2 lbs ae 2,4-D/acre) per application.
6. Observe a minimum of 30 days between applications.

ORNAMENTAL AND RECREATIONAL LAWNS AND TURF

Quali-Pro 3-D is for use on Ornamental Turf Lawns (Residential, Industrial, and Institutional), Parks, Cemeteries, Athletic Fields, and Golf Courses (Fairways, Aprons, Tees, and Roughs), and similar turf areas. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including but not limited to hospitals, nursing homes, schools, museums, libraries, sports facilities, golf courses, and office buildings. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings. Residential/domestic sites are defined as areas associated with the household or home life including but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

An adjuvant may be included in the spray solution.

Application is limited to 2 applications per year.

Use a maximum of 4.0 pts Quali-Pro 3-D/acre (1.2 lbs ae 2,4-D/acre) per application.

The maximum seasonal rate is 8.0 pts Quali-Pro 3-D/acre (2.4 lbs ae 2,4-D/acre) excluding spot treatments.

USE SITE	RATES	COMMENTS
Cool-Season Grasses such as Bentgrass Bluegrass Fescue Rye	3 to 4 pints per acre (1.1 to 1.5 fl oz/1000 sq ft) (.9 to 1.2 lb ae/acre)	For normal application, use 40 to 80 gallons of water per acre (1 to 2 gallons of water per 1000 sq ft). For professional lawn maintenance, use 1 to 5 gallons of water per 1000 sq ft. Higher water volumes may be used when tank mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.
Warm-Season Grasses such as St. Augustine (dormant) Bahia Bermuda Centipede (dormant) Zoysia	2 to 2.5 pints per acre (0.75 to 0.9 fl oz/1000 sq ft) (.6 to .8 lb ae/acre)	Use 40 to 200 gallons of water per acre (1 to 5 gallons of water per 1000 sq ft). Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm-season grasses. NOTE: If Bermudagrass is dormant, up to 4 pints per acre may be used (1.5 fl oz/1000 sq ft). However, some hybrid Bermudagrasses may be sensitive to this product. Contact your local Extension Service Weed Control Specialist.

NOTE: Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses, or injury may result. Large volumes of spray water (i.e., one fluid ounce in 5 gallons of water per 1500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet wetting each weed surface.

NON-TURF AREAS

Roadsides (including aprons and guardrails), rights-of-way, and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 2.6 to 6.5 pints (.8 to 2.0 lb ae 2,4-D) of Quali-Pro 3-D per 50 to

300 gallons of water. This mixture will cover 43,500 square feet. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of Quali-Pro 3-D per gallon of water and spray to thoroughly wet all foliage. **Postemergence (annual and perennial weeds) applications:** Application is limited to 2 applications per year. Use a maximum of 2.0 lbs ae 2,4-D or 6.5 pints of Quali-Pro 3-D per acre per application. Wait a minimum of 30 days between applications.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems, and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 6.0 to 13.0 pints of Quali-Pro 3-D to 100 gallons of water applying 200 to 600 gallons of spray mixture per 43,500 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying. **Postemergence applications (woody plants):** Limited to 1 application per year. Use a maximum of 13.0 pints Quali-Pro 3-D (4.0 lbs ae 2,4-D) per acre per year. Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use or for commercial seed production or for research purposes.

Use Precautions: Do not apply this product to St. Augustinegrass unless it is in the dormant stage. Care should be taken to avoid overdosing bentgrass, as injury may result. Some hybrid bermudagrass may be sensitive to this product. Contact your local Extension Service Weed Control Specialist for sensitive grasses in your area. Large volumes of spray water (i.e., one fluid ounce in 5 gallons water per 1500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: KEEP FROM FREEZING. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling. Do not store under conditions which might adversely affect the container or its ability to function properly.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applica-

ble law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro is a registered trademark of Makhteshim Agan of North America, Inc.

Compadre is a registered trademark of Loveland Products, Inc.